

10/561, 944

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1626KAS

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 "Ask CAS" for self-help around the clock  
NEWS 3 OCT 23 The Derwent World Patents Index suite of databases on STN  
has been enhanced and reloaded  
NEWS 4 OCT 30 CHEMLIST enhanced with new search and display field  
NEWS 5 NOV 03 JAPIO enhanced with IPC 8 features and functionality  
NEWS 6 NOV 10 CA/CAPLUS F-Term thesaurus enhanced  
NEWS 7 NOV 10 STN Express with Discover! free maintenance release Version  
8.01c now available  
NEWS 8 NOV 20 CA/CAPLUS to MARPAT accession number crossover limit increased  
to 50,000  
NEWS 9 DEC 01 CAS REGISTRY updated with new ambiguity codes  
NEWS 10 DEC 11 CAS REGISTRY chemical nomenclature enhanced  
NEWS 11 DEC 14 WPIDS/WPINDEX/WPIX manual codes updated  
NEWS 12 DEC 14 GBFULL and FRFULL enhanced with IPC 8 features and  
functionality  
NEWS 13 DEC 18 CA/CAPLUS pre-1967 chemical substance index entries enhanced  
with preparation role  
NEWS 14 DEC 18 CA/CAPLUS patent kind codes updated  
NEWS 15 DEC 18 MARPAT to CA/CAPLUS accession number crossover limit increased  
to 50,000  
NEWS 16 DEC 18 MEDLINE updated in preparation for 2007 reload  
NEWS 17 DEC 27 CA/CAPLUS enhanced with more pre-1907 records  
NEWS 18 JAN 08 CHEMLIST enhanced with New Zealand Inventory of Chemicals  
NEWS 19 JAN 16 CA/CAPLUS Company Name Thesaurus enhanced and reloaded  
NEWS 20 JAN 16 IPC version 2007.01 thesaurus available on STN  
NEWS 21 JAN 16 WPIDS/WPINDEX/WPIX enhanced with IPC 8 reclassification data  
NEWS 22 JAN 22 CA/CAPLUS updated with revised CAS roles  
NEWS 23 JAN 22 CA/CAPLUS enhanced with patent applications from India  
NEWS 24 JAN 29 PHAR reloaded with new search and display fields  
NEWS 25 JAN 29 CAS Registry Number crossover limit increased to 300,000 in  
multiple databases  
NEWS 26 FEB 13 CASREACT coverage to be extended  
NEWS 27 Feb 15 PATDPASPC enhanced with Drug Approval numbers  
NEWS 28 Feb 15 RUSSIAPAT enhanced with pre-1994 records  
  
NEWS EXPRESS NOVEMBER 10 CURRENT WINDOWS VERSION IS V8.01c, CURRENT  
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),  
AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.  
  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS LOGIN Welcome Banner and News Items  
NEWS IPC8 For general information regarding STN implementation of IPC 8  
NEWS X25 X.25 communication option no longer available

SAEED

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 11:27:51 ON 19 FEB 2007

=> FILE REG

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'REGISTRY' ENTERED AT 11:28:00 ON 19 FEB 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 18 FEB 2007 HIGHEST RN 921759-52-6

DICTIONARY FILE UPDATES: 18 FEB 2007 HIGHEST RN 921759-52-6

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 30, 2006


Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10439263.str



chain nodes :

7 8 10 11 12 13 17 18 19 21 27 28 29 30 31

ring nodes :

1 2 3 4 5 6 9 14 15 16 20 22 23 24 25 26

chain bonds :

4-7 7-8 7-10 8-9 8-13 10-11 11-12 12-31 16-17 17-18 17-21 18-19 19-20

24-27 27-28 27-29 29-30

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 9-14 9-16 14-15 15-16 20-22 20-26 22-23 23-24

24-25 25-26

exact/norm bonds :

7-10 8-9 8-13 9-14 9-16 10-11 12-31 14-15 15-16 17-18 17-21 18-19 27-28

27-29 29-30

exact bonds :

1-2 1-6 2-3 3-4 4-5 4-7 5-6 7-8 11-12 16-17 19-20 24-27

normalized bonds :

20-22 20-26 22-23 23-24 24-25 25-26

isolated ring systems :

containing 1 : 9 : 20 :

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS 8:CLASS 9:Atom 10:CLASS

11:CLASS 12:CLASS 13:CLASS 14:Atom 15:Atom 16:Atom 17:CLASS 18:CLASS

19:CLASS 20:Atom 21:CLASS 22:Atom 23:Atom 24:Atom 25:Atom 26:Atom 27:CLASS


28:CLASS 29:CLASS 30:CLASS 31:CLASS

L1 STRUCTURE UPLOADED

=> D

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> S L1

SAMPLE SEARCH INITIATED 11:28:21 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 32 TO ITERATE

100.0% PROCESSED 32 ITERATIONS

2 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
BATCH \*\*COMPLETE\*\*


PROJECTED ITERATIONS: 301 TO 979

PROJECTED ANSWERS: 2 TO 124

L2 2 SEA SSS SAM L1

=>

Uploading C:\Program Files\Stnexp\Queries\104392631.str



chain nodes :

SAEED

7 8 10 11 12 13 17 18 19 21 27 28 29 30 31 33  
ring nodes :  
1 2 3 4 5 6 9 14 15 16 20 22 23 24 25 26  
chain bonds :  
4-7 7-8 7-10 8-9 8-13 10-11 11-12 12-31 12-33 16-17 17-18 17-21 18-19  
19-20 24-27 27-28 27-29 29-30  
ring bonds :  
1-2 1-6 2-3 3-4 4-5 5-6 9-14 9-16 14-15 15-16 20-22 20-26 22-23 23-24  
24-25 25-26  
exact/norm bonds :  
7-10 8-9 8-13 9-14 9-16 10-11 14-15 15-16 17-18 17-21 18-19 27-28 27-29  
29-30  
exact bonds :  
1-2 1-6 2-3 3-4 4-5 4-7 5-6 7-8 11-12 16-17 19-20 24-27  
normalized bonds :  
12-31 12-33 20-22 20-26 22-23 23-24 24-25 25-26  
isolated ring systems :  
containing 1 : 9 : 20 :

## Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS 8:CLASS 9:Atom 10:CLASS  
11:CLASS 12:CLASS 13:CLASS 14:Atom 15:Atom 16:Atom 17:CLASS 18:CLASS  
19:CLASS 20:Atom 21:CLASS 22:Atom 23:Atom 24:Atom 25:Atom 26:Atom 27:CLASS  
28:CLASS 29:CLASS 30:CLASS 31:CLASS 33:CLASS

L3 STRUCTURE UPLOADED

=&gt; D

L3 HAS NO ANSWERS

L3 STR

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*

Structure attributes must be viewed using STN Express query preparation.

=&gt; S L3

SAMPLE SEARCH INITIATED 11:30:29 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 32 TO ITERATE

100.0% PROCESSED 32 ITERATIONS

0 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*

BATCH \*\*COMPLETE\*\*

PROJECTED ITERATIONS: 301 TO 979

PROJECTED ANSWERS: 0 TO 0

L4 0 SEA SSS SAM L3

=&gt; S L3 FULL

FULL SEARCH INITIATED 11:30:36 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 644 TO ITERATE

100.0% PROCESSED 644 ITERATIONS

1 ANSWERS

SEARCH TIME: 00.00.01

SAEED

L5 1 SEA SSS FUL L3

=> FILE CAPLUS  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
173.45	173.66

FULL ESTIMATED COST

FILE 'CAPLUS' ENTERED AT 11:30:41 ON 19 FEB 2007  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 19 Feb 2007 VOL 146 ISS 9  
FILE LAST UPDATED: 18 Feb 2007 (20070218/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> S L5  
L6 13 L5

=> D IBIB ABS HITSTR TOT

L6 ANSWER 1 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:460730 CAPLUS

DOCUMENT NUMBER: 145:201989

TITLE: Influence of erythromycin on the pharmacokinetics of ximelagatran may involve inhibition of P-glycoprotein-mediated excretion

AUTHOR(S): Eriksson, Ulf G.; Dorani, Hassan; Karlsson, Johan; Fritsch, Holger; Hoffmann, Kurt-Jürgen; Olsson, Lis; Sarich, Troy C.; Wall, Ulrike; Schutzer, Kaja-Marie; AstraZeneca R and D, Moelndal, Swed.

CORPORATE SOURCE: Drug Metabolism and Disposition (2006), 34(5), 775-782

PUBLISHER: American Society for Pharmacology and Experimental Therapeutics

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A pharmacokinetic interaction between erythromycin and ximelagatran, an oral direct thrombin inhibitor, was demonstrated in this study in healthy volunteers. To investigate possible interaction mechanisms, the effects of erythromycin on active transport mediated by P-glycoprotein (P-gp) in vitro in Caco-2 and P-gp-over-expressing Madin-Darby canine kidney-human multidrug resistance-1 cell preps. and on biliary excretion of melagatran in rats were studied. In healthy volunteers (seven males and nine females; mean age 24 years) receiving a single dose of ximelagatran 36 mg on day 1, erythromycin 500 mg t.i.d. on days 2 to 5, and a single dose of ximelagatran 36 mg plus erythromycin 500 mg on day 6, the least-squares mean ests. (90% confidence intervals) for the ratio of ximelagatran with erythromycin to ximelagatran given alone were 1.82 (1.64-2.01) for the area under the concentration-time curve and 1.74 (1.52-2.00) for the maximum plasma concentration of melagatran, the active form of ximelagatran. Neither the slope nor the intercept of the melagatran plasma concentration-effect relationship for activated partial thromboplastin time statistically significantly differed as a function of whether or not erythromycin was administered with ximelagatran. Ximelagatran was well tolerated regardless of whether it was administered with erythromycin. Erythromycin inhibited P-gp-mediated transport of both ximelagatran and melagatran in vitro and decreased the biliary excretion of melagatran in the rat. These results indicate that the mechanism of the pharmacokinetic interaction between oral ximelagatran and erythromycin may involve inhibition of transport proteins, possibly P-gp, resulting in decreased melagatran biliary excretion and increased bioavailability of melagatran.

IT 192939-72-3  
RL: BSU (Biological study, unclassified); FMU (Formation, unclassified); BIOL (Biological study); FORM (Formation, nonpreparative)  
(erythromycin effect on ximelagatran pharmacokinetics: P-gp mediation)

RN 192939-72-3 CAPLUS

CN Glycine, N-[(1R)-1-cyclohexyl-2-[(2S)-2-[[[4-[(hydroxyamino)iminomethyl]phenyl]methyl]amino]carbonyl]-1-azetidinyl]-2-oxoethyl]- (9CI) (CA INDEX NAME)

L6 ANSWER 2 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:498725 CAPLUS

DOCUMENT NUMBER: 143:126331

TITLE: The oral direct thrombin inhibitor, ximelagatran, an alternative for anticoagulant treatment during the puerperium and lactation

AUTHOR(S): Hellgren, M.; Johansson, S.; Eriksson, U. G.; Wahlender, K.

CORPORATE SOURCE: Department of Antenatal Care, Primary Health Care South Bohuslän and Institute for the Health of Women and Children, University of Goeteborg, Swed.

SOURCE: BJOG (2005), 112(5), 579-583

CODEN: BIOGPO; ISSN: 1470-0328

PUBLISHER: Blackwell Publishing Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Objective: To determine the excretion of the oral direct thrombin inhibitor (oral DTI), ximelagatran, and its active form, melagatran, in human milk, and to thus evaluate the potential exposure of breastfed infants to melagatran. Design: An open, single dose, single center study. Setting: Department of Antenatal Care, Primary Health Care South Bohuslän and Institute for the Health of Women and Children, Goeteborg University, Sweden. Sample: Seven healthy Caucasian breastfeeding women who were at least two months postpartum were studied. Methods: The concns. of ximelagatran, its two intermediates, and melagatran were determined using liquid

chromatog.-mass spectrometry, with the limit of quantification of 2 nmol L-1 for human milk and 10 nmol L-1 for plasma concns. Main outcome measures: Concns. of ximelagatran, its intermediates and melagatran were measured in breast milk over 72 h, and in plasma over 12 h, after a

single oral 36 mg dose of ximelagatran. Results: Neither ximelagatran nor its intermediates were detected in human breast milk. Only trace amts. of melagatran were detected. The mean cumulative amount of melagatran excreted

into breast milk over the 72-h period after dosing with oral ximelagatran was 0.00091% of the administered dose of ximelagatran. Ximelagatran was well tolerated, with no clin. relevant changes in laboratory variables or vitals.

signs. Conclusions: Trace levels of melagatran are excreted in human breast milk following administration of the oral DTI ximelagatran. The exposure of breastfed infants to melagatran appears to be low and is therefore unlikely to be of clin. concern.

IT 192939-72-3  
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(ximelagatran intermediate, hydroxy-melagatran was not excreted in breast milk of Caucasian women after oral DTI, ximelagatran administration)


RN 192939-72-3 CAPLUS

CN Glycine, N-[(1R)-1-cyclohexyl-2-[(2S)-2-[[[4-[(hydroxyamino)iminomethyl]phenyl]methyl]amino]carbonyl]-1-azetidinyl]-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L6 ANSWER 1 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)


Absolute stereochemistry.



REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE

FORMAT

L6 ANSWER 2 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)




REFERENCE COUNT: 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE

FORMAT

L6 ANSWER 3 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN  
 ACCESSION NUMBER: 2005:314009 CAPLUS  
 DOCUMENT NUMBER: 143:381615  
 TITLE: Characterization and partial purification of the rat and human enzyme systems active in the reduction of N-hydroxymelagatran and benzamidoxime  
 AUTHOR(S): Andersson, Susanne; Hofmann, Yvonne; Nordling, Asa; Li, Xue-qing; Nivelius, Sabina; Andersson, Tommy B.; Ingelman-Sundberg, Magnus; Johansson, Inger  
 CORPORATE SOURCE: Division of Molecular Toxicology, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Swed.  
 SOURCE: Drug Metabolism and Disposition (2005), 33(4), 570-578  
 CODEN: DMSAI; ISSN: 0090-9556  
 PUBLISHER: American Society for Pharmacology and Experimental Therapeutics  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB The enzymic basis for intracellular reduction of N-hydroxylated amidines to their corresponding amidines, and hydroxylamines to their corresponding amines, is unknown. The hydroxylated amidines can be used as prodrug moieties, and an understanding of the enzyme system active in the reduction can contribute to more efficient drug development. In this study, we examined the properties of this enzyme system using benzamidoxime and N-hydroxymelagatran as substrates. In rats and humans, the hepatic enzyme system was localized in mitochondria as well as in microsomes, using preferably NADH as cofactor. Potassium cyanide, N-methylhydroxylamine, p-hydroxymercuribenzoate, and desferrioxamine were efficient inhibitors, whereas typical cytochrome P 450 inhibitors were ineffective. In rats, the highest specific activity was found in liver, adipose tissue, and kidneys, whereas in humans, the specific activity in the preps. of adipose tissue examined was lower. A sex difference was observed in rat liver, where 4-fold higher activity was seen in microsomes from female rats. No gender differences were present in any other tissue investigated. Partial purification of the hepatic system was achieved using polyethylene glycol fractionation followed by Octyl Sepharose chromatog. at low detergent concns., whereas the enzyme was denatured after complete solubilization. The unique appearance of the enzyme activity in adipose tissue, together with the cyanide sensitivity and the failure of typical P 450 inhibitors to impede the reaction, indicates that the enzyme system active in reduction of benzamidoxime and N-hydroxymelagatran formation is not of cytochrome P 450 origin, but likely consists of an NADH-dependent electron transfer chain with a cyanide-sensitive protein as the terminal component.  
 IT 192939-72-3, Melagatran hydroxyamidine  
 RL: BSU (Biological study, unclassified); BIOL (Biological study) (characterization and partial purification of rat and human enzyme systems active in reduction of N-hydroxymelagatran and benzamidoxime)  
 RN 192939-72-3 CAPLUS

L6 ANSWER 3 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)  
 CN Glycine, N-[(1R)-1-cyclohexyl-2-[(2S)-2-[[[4-[(hydroxyamino)iminomethyl]phenyl]methyl]amino]carbonyl]-1-azetidinyl]-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

L6 ANSWER 4 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN  
 ACCESSION NUMBER: 2004:1156571 CAPLUS  
 DOCUMENT NUMBER: 142:56674  
 TITLE: New process for the production of melagatran  
 INVENTOR(S): Grehn, Marcus; Musil, Tibor; Sjoegren, Magnus  
 PATENT ASSIGNEE(S): AstraZeneca Ab, Swed.  
 SOURCE: PCT Int. Appl., 16 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004/113364	A1	2004-12-29	WO 2004-SE1016	2004-06-23
W:				
AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HK, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RN:				
BW, GH, GM, KE, LS, MM, MY, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2004249658	A1	2004-12-29	AU 2004-249658	2004-06-23
CA 2528930	A1	2004-12-29	CA 2004-2528930	2004-06-23
EP 1641814	A1	2006-04-05	EP 2004-749054	2004-06-23
R:				
AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, SI, LT, LV, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, HR				
CN 1809585	A	2006-07-26	CN 2004-80017545	2004-06-23
BR 2004011769	A	2006-08-08	BR 2004-11769	2004-06-23
NO 200505925	A	2006-01-19	NO 2005-5925	2005-12-13
US 2006178312	A1	2006-08-10	US 2005-561944	2005-12-21
PRIORITY APPLN. INFO.:			SE 2003-1879	A 2003-06-25
			WO 2004-SE1016	W 2004-06-23

OTHER SOURCE(S): CASREACT 142:56674; MARPAT 142:56674  
 AB A process for the production of melagatran [HO2CCH2-(R)Cgl-(S)Aze-Pab-H, where Cgl is cyclohexylglycyl, Aze is azetidine-2-carbonyl, and Pab is p-amidinobenzylamino] comprises the hydrolysis of an N-hydroxymelagatran alkyl or benzyl ester and reduction of the intermediate N-hydroxymelagatran. Thus, ximelagatran in ethanolic NaOH solution was stirred for four hours at 20-25°C to afford 90 weight % N-hydroxymelagatran, which was hydrogenated over 5% Pd/C to afford melagatran.  
 IT 192939-72-3P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) (production of melagatran from N-hydroxymelagatran esters)  
 RN 192939-72-3 CAPLUS  
 CN Glycine, N-[(1R)-1-cyclohexyl-2-[(2S)-2-[[[4-[(hydroxyamino)iminomethyl]phenyl]methyl]amino]carbonyl]-1-azetidinyl]-2-oxoethyl]- (9CI) (CA INDEX NAME)

L6 ANSWER 4 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)  
 Absolute stereochemistry.



REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

Applicat




L6 ANSWER 5 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN  
 ACCESSION NUMBER: 2004:104716 CAPLUS  
 DOCUMENT NUMBER: 142:62413  
 TITLE: Bioequivalence of ximelagatran, an oral direct thrombin inhibitor, as whole or crushed tablets or dissolved formulation  
 AUTHOR(S): Schuetzer, Kajs-Marie; Wall, Ulrika; Loennerstedt, Carina; Ohlsson, Lis; Teng, Renli; Sarich, Troy C.; Eriksson, Ulf G.  
 CORPORATE SOURCE: R+D Moelndal, AstraZeneca, Moelndal, Swed.  
 SOURCE: Current Medical Research and Opinion (2004), 20(3), 325-331  
 CODEN: CMROCK; ISSN: 0300-7995  
 PUBLISHER: LibraPharm Ltd.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB Objective: To investigate whether crushed or dissolved tablets of the oral direct thrombin inhibitor ximelagatran are bioequivalent to whole tablet administration. Ximelagatran is currently under development for the prevention and treatment of thromboembolic disorders. Research design and methods: This was an open-label, randomized, three-period, three-treatment crossover study in which 40 healthy volunteers (aged 20-33 yr) received a single 36-mg dose of ximelagatran administered in three different ways: I swallowed whole, II crushed, mixed with applesauce and ingested and III dissolved in water and administered via nasogastric tube. Results: The plasma concns. of ximelagatran, its intermediates and the active form melagatran were determined. Ximelagatran was rapidly absorbed and the bioavailability of melagatran was similar after the three different administrations, fulfilling the criteria for bioequivalence. The mean area under the plasma concentration-vs.-time curve (AUC) of melagatran was 1.6  $\mu\text{mol}\cdot\text{h/L}$  (ratio 1.01 for treatment II/I and 0.97 for treatment III/I), the mean peak concentration ( $C_{\text{max}}$ ) was 0.3  $\mu\text{mol/L}$  (ratio 1.04 for treatment II/I and 1.02 for treatment III/I) and the mean half-life ( $t_{1/2}$ ) was 2.8 h for all treatments. The time to  $C_{\text{max}}$  ( $t_{\text{max}}$ ) was 2.2 h for the whole tablet and approx. 0.5 h earlier when the tablet was crushed or dissolved (1.7-1.8 h), due to a more rapid absorption. The study drug was well tolerated as judged from the low incidence and type of adverse events reported. Conclusion: The present study showed that the pharmacokinetics (AUC and  $C_{\text{max}}$ ) of melagatran were not significantly altered whether ximelagatran was given orally as a crushed tablet mixed with applesauce or dissolved in water and given via nasogastric tube.  
 IT 192939-72-3  
 RL: BSU (Biological study, unclassified); BIOL (Biological study) (bioequivalence of ximelagatran as whole or crushed tablets or dissolved formulation)  
 RN 192939-72-3 CAPLUS  
 CN Glycine, N-[(1R)-1-cyclohexyl-2-[(2S)-2-[[[4-

L6 ANSWER 6 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN  
 ACCESSION NUMBER: 2003:701571 CAPLUS  
 DOCUMENT NUMBER: 140:138746  
 TITLE: Ximelagatran, an oral direct thrombin inhibitor, has a low potential for cytochrome P450-mediated drug-drug interactions  
 AUTHOR(S): Bredberg, Eva; Andersson, Tommy B.; Frison, Lars; Thuresson, Annelie; Johansson, Susanne; Eriksson-Lepkowska, Maria; Larsson, Marita; Eriksson, Ulf G.  
 CORPORATE SOURCE: AstraZeneca R and D, Moelndal, Swed.  
 SOURCE: Clinical Pharmacokinetics (2003), 42(8), 765-777  
 CODEN: CPKNDH; ISSN: 0312-5963  
 PUBLISHER: Adis International Ltd.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB Background: Ximelagatran is an oral direct thrombin inhibitor currently in clin. development for the prevention and treatment of thromboembolic disorders. After oral administration, ximelagatran is rapidly absorbed and extensively bioconverted, via two intermediates (ethyl-melagatran and hydroxy-melagatran), to its active form, melagatran. In vitro studies have shown no evidence for involvement of cytochrome P 450 (CYP) enzymes in either the bioactivation or the elimination of melagatran. Objective: To investigate the potential of ximelagatran, the intermediates ethyl-melagatran and hydroxy-melagatran, and melagatran to inhibit the CYP system in vitro and in vivo, and the influence of three CYP substrates on the pharmacokinetics of melagatran in vivo. Methods: The CYP inhibitory properties of ximelagatran, the intermediates and melagatran were tested in vitro by two different methods, using heterologously expressed enzymes or human liver microsomes. Diclofenac (CYP2C9), diazepam (CYP2C19) and nifedipine (CYP3A4) were chosen for coadministration with ximelagatran in healthy volunteers. Subjects received oral ximelagatran 24 mg and/or diclofenac 50 mg, a 10-min i.v. infusion of diazepam 0.1 mg/kg, or nifedipine 60 mg. The plasma pharmacokinetics of melagatran, diclofenac, diazepam, N-desmethyl-diazepam and nifedipine were determined when administered alone and in combination with ximelagatran. Results: No inhibition, or only minor inhibition, of CYP enzymes by ximelagatran, the intermediates or melagatran was shown in the in vitro studies, suggesting that ximelagatran would not cause CYP-mediated drug-drug interactions in vivo. This result was confirmed in the clin. studies. There were no statistically significant differences in the pharmacokinetics of diclofenac, diazepam and nifedipine on coadministration with ximelagatran. Moreover, there were no statistically significant differences in the pharmacokinetics of melagatran when ximelagatran was administered alone or in combination with diclofenac, diazepam or nifedipine. Conclusion: As ximelagatran did not exert a significant effect on the hepatic CYP isoenzymes responsible for the metabolism of diclofenac, diazepam and nifedipine, it is reasonable to expect that it would have no effect on the metabolism of other drugs metabolized by these isoenzymes. Furthermore, the pharmacokinetics of melagatran after oral administration of ximelagatran are not expected to be altered by inhibition or induction of CYP2C9, CYP2C19 or CYP3A4. Together, the in vitro and in vivo studies indicate

L6 ANSWER 5 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)  
 [(hydroxyamino)iminomethyl]phenyl)methyl]amino]carbonyl]-1-azetidiny]-2-oxoethyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L6 ANSWER 6 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)  
 that metabolic drug-drug interactions involving the major human CYP enzymes should not be expected with ximelagatran.  
 IT 192939-72-3  
 RL: PAC (Pharmacological activity); PKT (Pharmacokinetics); BIOL (Biological study) (oral direct thrombin inhibitor ximelagatran has low potential for cytochrome P 450-mediated drug-drug interactions in vitro and in vivo in humans)  
 RN 192939-72-3 CAPLUS  
 CN Glycine, N-[(1R)-1-cyclohexyl-2-[(2S)-2-[[[4-[(hydroxyamino)iminomethyl]phenyl)methyl]amino]carbonyl]-1-azetidiny]-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT


L6 ANSWER 7 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN  
 ACCESSION NUMBER: 2003:701570 CAPLUS  
 DOCUMENT NUMBER: 139:270230  
 TITLE: No influence of mild-to-moderate hepatic impairment on the pharmacokinetics and pharmacodynamics of ximelagatran, an oral direct thrombin inhibitor  
 AUTHOR(S): Wahlander, Karin; Eriksson-Lepkowska, Maria; Frison, Lars; Fager, Gunnar; Eriksson, Ulf G.  
 CORPORATE SOURCE: AstraZeneca R and D, Mölndal, Swed.  
 SOURCE: Clinical Pharmacokinetics (2003), 42(8), 755-764  
 CODEN: CPKNDH; ISSN: 0312-5963  
 PUBLISHER: Adis International Ltd.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB Background: The oral direct thrombin inhibitor ximelagatran is a new class of anticoagulant currently in clin. development for the prevention and treatment of thromboembolic diseases. After oral administration, ximelagatran is rapidly absorbed and bioconverted to its active form melagatran. Objective: To investigate the influence of mild-to-moderate hepatic impairment on the pharmacokinetic and pharmacodynamic properties of ximelagatran. Study design: Nonblinded, nonrandomised study. Participants: Twelve volunteers with mild-to-moderate hepatic impairment (classified as Child-Pugh A or B) and 12 age-, weight-, and sex-matched control volunteers with normal hepatic function. Methods: Volunteers received a single oral dose of ximelagatran 24mg. Plasma and urine samples were collected for pharmacokinetic and pharmacodynamic analyses. Results: The absorption and bioconversion of ximelagatran to melagatran were rapid in both groups. The maximum plasma concentration of melagatran (C<sub>max</sub>) was achieved 2-3 h after administration; the mean elimination half-life (t<sub>1/2z</sub>) was 3.6 h for hepatically impaired volunteers and 3.1 h for the control volunteers. The area under the plasma concentration-time curve (AUC) and C<sub>max</sub> of melagatran in volunteers with hepatic impairment were 11 and 25% lower than in control volunteers, resp. However, after correcting for the higher renal function (i.e. higher calculated creatinine clearance) in the hepatically impaired volunteers, the ratio of melagatran AUC for hepatically impaired/control volunteers was 0.98 (90% CI 0.80, 1.22), suggesting that mild-to-moderate hepatic impairment had no influence on the pharmacokinetics of ximelagatran. Melagatran was the predominant compound in urine, accounting for 13-14% of the ximelagatran dose. Renal clearance of melagatran was 13% higher in hepatically impaired than in control volunteers. There were no significant differences between the two groups in the concentration-response relation between plasma melagatran concentration and activated partial thromboplastin time (APTT). Baseline prothrombin time (PT) was slightly longer in the hepatically impaired patients than in the control volunteers, probably reflecting a slight decrease in the activity of coagulation factors. However, when concns. of melagatran were

L6 ANSWER 8 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN  
 ACCESSION NUMBER: 2003:497014 CAPLUS  
 DOCUMENT NUMBER: 139:390673  
 TITLE: No influence of obesity on the pharmacokinetics and pharmacodynamics of melagatran, the active form of the oral direct thrombin inhibitor ximelagatran  
 AUTHOR(S): Sarich, Troy C.; Teng, Renli; Peters, Gary R.; Wollbrett, Maria; Homolka, Robert; Svensson, Mia; Eriksson, Ulf G.  
 CORPORATE SOURCE: AstraZeneca LP, Wilmington, DE, USA  
 SOURCE: Clinical Pharmacokinetics (2003), 42(5), 485-492  
 CODEN: CPKNDH; ISSN: 0312-5963  
 PUBLISHER: Adis International Ltd.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB Background: Ximelagatran, an oral direct thrombin inhibitor, is currently in clin. development for the prevention and treatment of thromboembolic diseases. Following oral administration, ximelagatran undergoes rapid bioconversion to its active form, melagatran, via two minor intermediates. Obesity, defined as body mass index (BMI) >30 kg/m<sup>2</sup>, is a recognized risk factor for thrombosis. There is potential for differences in the pharmacokinetics and pharmacodynamics of drugs administered to obese vs. non-obese patients, and some drugs may require alternative administration strategies in obese patients. Objective: To investigate the effect of obesity on the pharmacokinetics and pharmacodynamics of melagatran after oral administration of ximelagatran. Design and participants: This was an open-label, single-dose, group-matched study in which obese subjects (BMI 32-39 kg/m<sup>2</sup>; six male and six female; age 21-40 yr) were matched by sex and age (±2 yr) with non-obese subjects (BMI 21-26 kg/m<sup>2</sup>; six male and six female; aged 21-39 yr). Each subject received a single oral dose of ximelagatran 24mg. Blood samples for determination of plasma concns. of melagatran and activated partial thromboplastin times (APTT; a marker of melagatran pharmacodynamics) were collected up to 12 h after administration. Results: There were no statistically significant differences in the pharmacokinetic properties of melagatran between obese and non-obese subjects. Values of area under the melagatran plasma concentration-time curve, maximum plasma concentration, (C<sub>max</sub>), time at which C<sub>max</sub> occurred and terminal elimination half-life were approx. 1 μmol = h/L, 0.2 μmol/L, 2 h and 3 h in obese and non-obese subjects, resp. In addition, there was no statistically significant difference between the obese and non-obese subjects in the amount of ximelagatran, melagatran or the minor intermediates ethyl-melagatran and melagatran hydroxyamide excreted in urine. When relating the prolongation of APTT ratio to the square root of plasma concentration of melagatran and obesity status (no/yes), no statistically significant interaction between plasma concentration and obesity status was observed. Ximelagatran was well tolerated in both obese and non-obese subjects, and no bleeding events or serious adverse events occurred. Conclusion: No differences in the pharmacokinetics or pharmacodynamics of melagatran were detected between obese and non-obese subjects after oral administration of ximelagatran, suggesting that dose adjustment of ximelagatran in obesity (BMI up to 39 kg/m<sup>2</sup>) is not necessary.  
 IT 192939-72-3, Melagatran hydroxyamide

SAEED

L6 ANSWER 7 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)  
 at their peak, the increase in PT from baseline values was the same in both groups. Capillary bleeding time was measured in the hepatically impaired patients only, and was not increased by ximelagatran. Ximelagatran was well tolerated in both groups. Conclusion: There were no differences in the pharmacokinetic or pharmacodynamic properties of melagatran following oral administration of ximelagatran between the hepatically impaired and control volunteers. These findings suggest that dose adjustment for patients with mild-to-moderate impairment of hepatic function is not necessary.  
 IT 192939-72-3, Melagatran hydroxyamide  
 RL: ADV (Adverse effect, including toxicity); BSU (Biological study, unclassified); PAC (Pharmacological activity); PKT (Pharmacokinetics); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (no influence of mild-to-moderate hepatic impairment on pharmacokinetics and pharmacodynamics of ximelagatran (Exanta))  
 RN 192939-72-3 CAPLUS  
 CN Glycine, N-[(1R)-1-cyclohexyl-2-[(2S)-2-[[[4-[(hydroxyamino)iminomethyl]phenyl]methyl]amino]carbonyl]-1-azetidinyl]-2-oxoethyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

L6 ANSWER 8 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)  
 RL: PKT (Pharmacokinetics); BIOL (Biological study) (influence of obesity on the pharmacokinetics and pharmacodynamics of melagatran)  
 RN 192939-72-3 CAPLUS  
 CN Glycine, N-[(1R)-1-cyclohexyl-2-[(2S)-2-[[[4-[(hydroxyamino)iminomethyl]phenyl]methyl]amino]carbonyl]-1-azetidinyl]-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

L6 ANSWER 9 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER:

2003:49703 CAPLUS

DOCUMENT NUMBER:

139:39072

TITLE:

No influence of ethnic origin on the pharmacokinetics and pharmacodynamics of melagatran following oral administration of ximelagatran, a novel oral direct thrombin inhibitor, to healthy male volunteers  
Johansson, Linda C.; Andersson, Magnus; Fager,

AUTHOR(S):

Gunnar;

CORPORATE SOURCE:

Gustafsson, David; Eriksson, Ulf G.

SOURCE:

AstraZeneca R&amp;D, Moelndal, Sweden.

PUBLISHER:

Clinical Pharmacokinetics (2003), 42(5), 475-484

DOCUMENT TYPE:

CODEN: CPKNDH; ISSN: 0312-5963

LANGUAGE:

Adis International Ltd.

AB Objective:

Journal

pharmacokinetic

English

and pharmacodynamic properties of melagatran after oral administration of ximelagatran, a novel oral direct thrombin inhibitor. Study design: This was an open-label, non-randomized study with a single study session.

Subjects:

Thirty-six young healthy male subjects living in France were divided equally according to their ethnic origin (African, Asian and Caucasian). Methods: All subjects received a single 50mg oral dose of ximelagatran in solution. Blood and urine samples for pharmacokinetic evaluation were collected up to 12 and 24 h after administration, resp.

Blood samples were also collected to determine the activated partial thromboplastin time (APTT), an ex vivo coagulation time measurement used to demonstrate inhibition of thrombin, up to 24 h after administration.

Results:

The absorption of ximelagatran, and its bioconversion to melagatran, was rapid in all three ethnic groups. The metabolite pattern in plasma and urine was similar in all groups, with melagatran being the dominant compound. For ximelagatran, the mean area under the plasma concentration-time curve (AUC) was similar in the three groups, suggesting that

there was no difference in the extent to which ximelagatran was absorbed. Melagatran AUC was higher in the Asian subjects, with a mean Asian/Caucasian ratio (95% CI) of 1.23 (1.04, 1.45). This was presumably because of their lower bodyweight, which is correlated to lower renal function. Following normalization for bodyweight, there were no statistically significant differences between the three ethnic groups. This finding suggests that renal elimination was lower for Asian subjects,

whereas there were no differences in the conversion of ximelagatran to melagatran. The interindividual variability of melagatran AUC was low (coefficient of variation 19-26%), and the mean bioavailability of melagatran, estimated using a mean value for melagatran clearance obtained from Caucasian subjects in a previous study, was approx. 20% in all groups (range of mean values 19-23%). APTT increased nonlinearly with increasing melagatran plasma concentration, and no difference in the concentration-response relationship was observed between the groups. Conclusion: After oral administration of ximelagatran, the pharmacokinetic and pharmacodynamic properties of

ximelagatran, an amidoxime and ester prodrug

Clement, Bernd; Lopian, Katrin

Pharmaceutical Institute, Christian-Albrechts-University of Kiel, Kiel, Germany

Drug Metabolism and Disposition (2003), 31(5), 645-651

CODEN: DMSDAI; ISSN: 0090-9556

American Society for Pharmacology and Experimental Therapeutics

Journal

English

N-Hydroxylated amidines (amidoximes) can be used as prodrugs of amidines. The prodrug principle was developed in our laboratory for pentamidine and had been applied to several other drug candidates. One of these compds. is melagatran, a novel, synthetic, low mol. weight, direct thrombin inhibitor.

To increase the poor oral bioavailability due to its strong basic amidine functionality selected to fit the arginine side pocket of thrombin, the less basic N-hydroxylated amidine was used in addition to an Et ester-protecting residue. The objective of this investigation was to study the reduction and the hydrolytic metabolism of ximelagatran via two mono-prodrugs (N-hydroxy-melagatran and ethyl-melagatran) to melagatran

in vitro expts. New high-performance liquid chromatog. methods were developed to analyze all four compds. The biotransformation of ximelagatran to melagatran involving the reduction of the amidoxime function

and the ester cleavage could be demonstrated in vitro by microsomes and mitochondria from liver and kidney of pig and human, and the kinetic parameters were determined. So far, one enzyme system capable of reducing N-hydroxylated structures has been identified in pig liver microsomes, consisting of cytochrome b5, NADH-cytochrome b5 reductase, and a P 450 isoenzyme of the subfamily 2D. This enzyme system also reduces ximelagatran and N-hydroxy-melagatran. The participation of recombinant human CYP1A2, 2A6, 2C9, 2C19, 2D6, and 3A4 with cytochrome b5 and b5 reductase in the reduction can be excluded. In summary, ximelagatran and N-hydroxy-melagatran are easily reduced by several enzyme systems located in microsomes and mitochondria of different organs.

IT 192939-72-3, N-Hydroxymelagatran

RL: PKT (Pharmacokinetics); BIOL (Biological study)

(in vitro biotransformation of ximelagatran, an amidoxime and ester prodrug, and role of microsomal and mitochondrial enzymes)

192939-72-3 CAPLUS

Glycine, N-[(1R)-1-cyclohexyl-2-[(2S)-2-[[[4-[(hydroxyamino)iminomethyl]phenyl]methyl]amino]carbonyl]-1-azetidiny]-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

SAEED

L6 ANSWER 9 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)

melagatran are independent of ethnic origin. The elimination of melagatran is correlated with renal function.

IT

192939-72-3, Melagatran hydroxyamidine

RL: PKT (Pharmacokinetics); BIOL (Biological study)


(influence of ethnic origin on the pharmacokinetics and pharmacodynamics of melagatran following oral administration of ximelagatran to healthy male volunteers)

RN

192939-72-3 CAPLUS

Glycine, N-[(1R)-1-cyclohexyl-2-[(2S)-2-[[[4-[(hydroxyamino)iminomethyl]phenyl]methyl]amino]carbonyl]-1-azetidiny]-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT:

21

THIS

THERE ARE 21 CITED REFERENCES AVAILABLE FOR

FORMAT

RECORD. ALL CITATIONS AVAILABLE IN THE RE

L6 ANSWER 10 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER:

2003:125363 CAPLUS

DOCUMENT NUMBER:

139:270214

TITLE:

Characterization of in vitro biotransformation of new, orally active, direct thrombin inhibitor

ximelagatran,

an amidoxime and ester prodrug

Clement, Bernd; Lopian, Katrin

Pharmaceutical Institute, Christian-Albrechts-University of Kiel, Kiel, Germany

Drug Metabolism and Disposition (2003), 31(5), 645-651

CODEN: DMSDAI; ISSN: 0090-9556

American Society for Pharmacology and Experimental Therapeutics

Journal

English

N-Hydroxylated amidines (amidoximes) can be used as prodrugs of amidines. The prodrug principle was developed in our laboratory for pentamidine and had been applied to several other drug candidates. One of these compds. is melagatran, a novel, synthetic, low mol. weight, direct thrombin inhibitor.

To increase the poor oral bioavailability due to its strong basic amidine functionality selected to fit the arginine side pocket of thrombin, the less basic N-hydroxylated amidine was used in addition to an Et ester-protecting residue. The objective of this investigation was to study the reduction and the hydrolytic metabolism of ximelagatran via two mono-prodrugs (N-hydroxy-melagatran and ethyl-melagatran) to melagatran

in vitro expts. New high-performance liquid chromatog. methods were developed to analyze all four compds. The biotransformation of ximelagatran to melagatran involving the reduction of the amidoxime function

and the ester cleavage could be demonstrated in vitro by microsomes and mitochondria from liver and kidney of pig and human, and the kinetic parameters were determined. So far, one enzyme system capable of reducing N-hydroxylated structures has been identified in pig liver microsomes, consisting of cytochrome b5, NADH-cytochrome b5 reductase, and a P 450 isoenzyme of the subfamily 2D. This enzyme system also reduces ximelagatran and N-hydroxy-melagatran. The participation of recombinant human CYP1A2, 2A6, 2C9, 2C19, 2D6, and 3A4 with cytochrome b5 and b5 reductase in the reduction can be excluded. In summary, ximelagatran and N-hydroxy-melagatran are easily reduced by several enzyme systems located in microsomes and mitochondria of different organs.

IT 192939-72-3, N-Hydroxymelagatran

RL: PKT (Pharmacokinetics); BIOL (Biological study)

(in vitro biotransformation of ximelagatran, an amidoxime and ester prodrug, and role of microsomal and mitochondrial enzymes)


192939-72-3 CAPLUS

Glycine, N-[(1R)-1-cyclohexyl-2-[(2S)-2-[[[4-[(hydroxyamino)iminomethyl]phenyl]methyl]amino]carbonyl]-1-azetidiny]-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

SAEED

L6 ANSWER 10 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)



REFERENCE COUNT:

36

THIS

THERE ARE 36 CITED REFERENCES AVAILABLE FOR

FORMAT


RECORD. ALL CITATIONS AVAILABLE IN THE RE

L6 ANSWER 11 OF 13 CAPLUS COPYRIGHT 2007 ACS ON STN  
 ACCESSION NUMBER: 2003:15659 CAPLUS  
 DOCUMENT NUMBER: 139:223644  
 TITLE: Absorption, distribution, metabolism, and excretion of  
 ximelagatran, an oral direct thrombin inhibitor, in rats, dogs, and humans  
 AUTHOR(S): Eriksson, Ulf G.; Bredberg, Ulf; Hoffmann, Kurt-Jürgen; Thuresson, Anneli; Gabriellsson, Margareth; Ericsson, Hans; Ahnoff, Martin; Gislén, Kristina; Fager, Gunnar; Gustafsson, David  
 CORPORATE SOURCE: AstraZeneca R and D Mölndal, Mölndal, S-431 83, Sweden  
 SOURCE: Drug Metabolism and Disposition (2003), 31(3), 294-305  
 CODEN: DMDSAI; ISSN: 0090-9556  
 PUBLISHER: American Society for Pharmacology and Experimental Therapeutics  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB The absorption, metabolism, and excretion of the oral direct thrombin inhibitor, ximelagatran, and its active form, melagatran, were investigated in rats, dogs, and healthy male human subjects after administration of oral and i.v. single doses. Ximelagatran was rapidly absorbed and metabolized following oral administration, with melagatran as the predominant compound in plasma. Two intermediates (ethyl-melagatran and OH-melagatran) that were subsequently metabolized to melagatran were also identified in plasma and were rapidly eliminated. Melagatran given i.v. had relatively low plasma clearance, small volume of distribution, and short elimination half-life. The oral absorption of melagatran was low and highly variable. It was primarily renally cleared, and the renal clearance agreed well with the glomerular filtration rate. Ximelagatran was extensively metabolized, and only trace amounts were renally excreted. Melagatran was the major compound in urine and feces after administration of ximelagatran. Appreciable quantities of ethyl-melagatran were also recovered in rat, dog, and human feces after oral administration, suggesting reduction of the hydroxyamidine group of ximelagatran in the gastrointestinal tract, as demonstrated when ximelagatran was incubated with feces homogenate. Polar metabolites in urine and feces (all species) accounted for a relatively small fraction of the dose. The bioavailability of melagatran following oral administration of ximelagatran was 5 to 10% in rats, 10 to 50% in dogs, and about 20% in humans, with low between-subject variation. The fraction of ximelagatran absorbed was at least 40 to 70% in all species. First-pass metabolism of ximelagatran with subsequent biliary excretion of the formed metabolites account for the lower bioavailability of melagatran.  
 IT 192939-72-3  
 RL: PKT (Pharmacokinetics); BIOL (Biological study)  
 (absorption, distribution, metabolism, and excretion of ximelagatran, an oral direct thrombin inhibitor, in rats, dogs, and humans)

L6 ANSWER 12 OF 13 CAPLUS COPYRIGHT 2007 ACS ON STN  
 ACCESSION NUMBER: 2003:99008 CAPLUS  
 DOCUMENT NUMBER: 139:62516  
 TITLE: Determination of ximelagatran, an oral direct thrombin  
 inhibitor, its active metabolite melagatran, and the intermediate metabolites, in biological samples by liquid chromatography-mass spectrometry  
 AUTHOR(S): Larsson, Marita; Ahnoff, Martin; Abrahamsson, Anna; Logren, Ulrika; Fakt, Christina; Ohlman, Irene; Persson, Bengt-Arne  
 CORPORATE SOURCE: DMPK and Bioanalytical Chemistry, AstraZeneca R and D Mölndal, Mölndal, S-431 83, Sweden  
 SOURCE: Journal of Chromatography, B: Analytical Technologies in the Biomedical and Life Sciences (2003), 783(2), 335-347  
 CODEN: JCBAAI; ISSN: 1570-0232  
 PUBLISHER: Elsevier Science B.V.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB Anal. methods for the determination of ximelagatran, an oral direct thrombin inhibitor, its active metabolite melagatran, and intermediate metabolites, melagatran hydroxyamidine and melagatran Et ester, in biol. samples by liquid chromatog. (LC) pos. electrospray ionization mass spectrometry (MS) using selected reaction monitoring are described. Isolation from human plasma was achieved by solid-phase extraction on octylsilica. Analytes and isotope-labeled internal stds. were separated by LC utilizing a C18 anal. column and a mobile phase comprising acetonitrile-4 mmol/L ammonium acetate (35:65, volume/volume) containing 0.1% formic acid, at a flow-rate of 0.75 mL/min. Absolute recovery was approx. 80% for ximelagatran, approx. 60% for melagatran Et ester, and >90% for melagatran and melagatran hydroxyamidine. Limit of quantification was 10 nmol/L, with a relative standard deviation <20% for each analyte and <5% above 100 nmol/L.  
 Procedures for the determination of these analytes in human urine and breast milk, plus whole blood from rat and mouse are also described.  
 IT 192939-72-3, Melagatran hydroxyamidine  
 RL: ANT (Analyte); ANST (Analytical study)  
 (ximelagatran and its active metabolite melagatran and intermediate metabolites determination in biol. samples by liquid chromatog.-mass spectrometry)  
 RN 192939-72-3 CAPLUS  
 CN Glycine, N-[(1R)-1-cyclohexyl-2-[(2S)-2-[[[4-[(hydroxyamino)iminomethyl]phenyl]methyl]amino]carbonyl]-1-azetidinyl]-2-oxoethyl]- (9CI) (CA INDEX NAME)  
 Absolute stereochemistry.


L6 ANSWER 11 OF 13 CAPLUS COPYRIGHT 2007 ACS ON STN (Continued)  
 RN 192939-72-3 CAPLUS  
 CN Glycine, N-[(1R)-1-cyclohexyl-2-[(2S)-2-[[[4-[(hydroxyamino)iminomethyl]phenyl]methyl]amino]carbonyl]-1-azetidinyl]-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

L6 ANSWER 12 OF 13 CAPLUS COPYRIGHT 2007 ACS ON STN (Continued)



REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

L6 ANSWER 13 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN  
 ACCESSION NUMBER: 1997:506597 CAPLUS  
 DOCUMENT NUMBER: 127:136080  
 TITLE: Preparation of peptide derivatives as prodrugs of thrombin inhibitors  
 INVENTOR(S): Antonsson, Thomas; Gustafsson, David; Hoffmann, Kurt-Jurgen; Nystrom, Jan-Erik; Sorensen, Henrik; Sellen, Mikael  
 PATENT ASSIGNEE(S): Astra Aktiebolag, Swed.  
 SOURCE: PCT Int. Appl., 94 pp.  
 CODEN: PIXX2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9723499	A1	19970703	WO 1996-SE1680	19961217
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
ZA 9610353	A	19970623	ZA 1996-10353	19961209
TW 541316	B	20030711	TW 1996-B5115209	19961209
TW 238827	B	20050901	TW 2002-91115525	19961209
CA 2238737	A1	19970703	CA 1996-2238737	19961217
AU 9712178	A	19970717	AU 1997-12178	19961217
AU 706350	B2	19990617		
EP 869966	A1	19981014	EP 1996-943446	19961217
EP 869966	B1	20050316		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
CN 1209139	A	19990224	CN 1996-180024	19961217
CN 1127510	B	20031112		
HU 9900115	A2	19990528	HU 1999-115	19961217
BR 9612148	A	19990713	BR 1996-12148	19961217
JP 2000504313	T	20000411	JP 1997-523571	19961217
JP 3282821	B2	20020520		
EP 995755	A1	20000426	EP 1999-120315	19961217
EP 995755	B1	20010816		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
NZ 324902	A	20000623	NZ 1996-324902	19961217
AT 204292	T	20010915	AT 1999-120315	19961217
RU 2176644	C2	20011210	RU 1998-111148	19961217
ES 2163916	T3	20020201	ES 1999-120315	19961217
PT 995755	T	20020228	PT 1999-120315	19961217
EE 4022	B1	20030415	EE 1998-187	19961217
PL 187468	B1	20040730	PL 1996-327569	19961217
NZ 504245	A	20041126	NZ 1996-504245	19961217
AT 291031	T	20050415	AT 1996-943446	19961217
EP 1533318	A1	20050525	EP 2005-116	19961217


L6 ANSWER 13 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO  
 PT 869966 T 20050729 PT 1996-943446 19961217  
 ES 2128283 T3 20050901 ES 1996-943446 19961217  
 US 5965692 A 19991012 US 1997-776231 19970131  
 NO 9802809 A 19980820 NO 1998-2809 19980618  
 HK 1016610 A1 20050610 HK 1999-101429 19990409  
 US 6262028 B1 20010717 US 1999-353644 19990715  
 JP 2001089498 A 20010403 JP 2000-220423 20000721  
 JP 3580535 B2 20041027  
 HK 1026214 A1 20020809 HK 2000-105419 20000830  
 US 2002142968 A1 20021003 US 2002-74008 20020214  
 US 2004162410 A1 20040819 US 2004-776245 20040212  
 GB 1995-26273 A 19951221

PRIORITY APPLN. INFO.:  
 SE 1996-556 A 19960215  
 EP 1996-943446 A3 19961217  
 JP 1997-523571 A3 19961217  
 WO 1996-SE1680 W 19961217  
 US 1997-776231 A1 19970131  
 SE 1997-4542 A 19971205  
 WO 1998-SE2191 A 19981201  
 US 1999-353644 A1 19990715  
 US 2000-708449 B1 20001109  
 US 2002-74008 B1 20020214

OTHER SOURCE(S): MARPAT 127:136080  
 AB Title compds. of formula R1O(O)C-CH2-(R)Cgl-Aze-Pab-R2 [wherein R1 = H, C1-10 alkyl, (un)substituted C1-3 alkylphenyl, A1C(O)N(R3)R4, A1C(O)OR3; (R)Cgl = (R)-cyclohexyl glycine; Aze = (S)-azetidine-2-carboxylic acid; Pab = 1-amidino-4-aminomethylbenzene; R2 (which replaces one of the hydrogen atoms in the amidino unit of Pab) = OH, OC(O)R5, C(O)OR6, C(O)OCH(R7)OC(O)R8; R3 and R4 are independently e.g., H, C1-6 alkyl, Ph, or together with the nitrogen atom represent pyrrolidinyl or piperidinyl; R5 = C1-17 alkyl, Ph, or 2-naphthyl (all of which are optionally substituted by C1-6 alkyl or halogen); R6 = (un)substituted 2-naphthyl, Ph, C1-3 alkylphenyl, C1-12 alkyl; R7 = H, C1-4 alkyl; R8 = e.g., 2-naphthyl, Ph, C1-6 alkoxy, (un)substituted C1-8 alkyl] or a pharmaceutically acceptable salt thereof, which are useful as prodrugs of inhibitors of trypsin-like proteases (no data), such as thrombin, and in particular in the treatment of conditions where inhibition of thrombin is required (e.g. thrombosis) or as anticoagulants, were prepared For example, EtO2C-CH2-(R)Cgl-Aze-Pab-COOCH2CH:CH2 was prepared via coupling of Me3CO2C-(R)Cgl-Aze-Pab-H with allyl chloroformate followed by Boc deprotection and coupling with Et bromoacetate. The title compds. were

L6 ANSWER 13 OF 13 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)  
 all found to exhibit oral and/or parenteral bioavailability in rats as the active inhibitor HO2C-CH2-(R)Cgl-Aze-Pab-H, either as the free acid and/or as one or more ester thereof.  
 IT 192939-72-3P  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)  
 (preparation of peptide derivs. as prodrugs of thrombin inhibitors)  
 RN 192939-72-3 CAPLUS  
 CN Glycine, N-[(1R)-1-cyclohexyl-2-[[[2S]-2-[[[4-(hydroxyamino)iminomethyl]phenyl]methyl]amino]carbonyl]-1-azetidinyl]-2-oxoethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



=> S L6 AND MELAGATRAN  
252 MELAGATRAN  
L7 12 L6 AND MELAGATRAN

=> LOGOFF

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

LOGOFF? (Y)/N/HOLD:Y

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
74.75	248.41

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
-10.14	-10.14

CA SUBSCRIBER PRICE

STN INTERNATIONAL LOGOFF AT 11:36:09 ON 19 FEB 2007